1) Alzheimer's Disease/Severe Dementia

2014

2019

Deterioration or loss of intellectual capacity as confirmed by clinical evaluation and imaging tests, arising from Alzheimer's Disease or irreversible organic disorders, resulting in significant reduction in mental and social functioning requiring the continuous supervision of the Life Assured.

This diagnosis must be supported by the clinical confirmation of an appropriate consultant and supported by Our appointed doctor.

The following are excluded:

- (a) Non-organic diseases such as neurosis and psychiatric illnesses; and
- (b) Alcohol related brain damage.

Deterioration or loss of cognitive function as confirmed by clinical evaluation and imaging tests, arising from Alzheimer's disease or irreversible organic disorders, resulting in significant reduction in mental and social functioning requiring the continuous supervision of the life assured.

This diagnosis must be supported by the clinical confirmation of an appropriate consultant and supported by the Company's appointed doctor.

The following are excluded:

- Non-organic diseases such as neurosis and psychiatric illnesses; and
- · Alcohol related brain damage.

No significant change

2) Angioplasty & Other Invasive Treatment for Coronary Artery 2014 2019 The actual undergoing of balloon The actual undergoing of balloon angioplasty or similar intra-arterial angioplasty or similar intra-arterial catheter procedure to correct a narrowing catheter procedure to correct a of minimum 60% stenosis, of one or more narrowing of minimum 60% stenosis, of major coronary arteries as shown by one or more major coronary arteries as angiographic evidence. The shown by angiographic evidence. The revascularisation must be considered revascularisation must be considered medically necessary by a consultant medically necessary by a consultant cardiologist. cardiologist. Coronary arteries herein refer to left main Coronary arteries herein refer to left stem, left anterior descending, circumflex main stem, left anterior descending, and right coronary artery. circumflex and right coronary artery. Payment under this condition is limited to 10% of the Sum Assured under this policy subject to a S\$25,000 maximum sum payable. This benefit is payable once only and shall be deducted from the amount of this Contract, thereby reducing the amount of the Sum Assured which may be payable herein.

Diagnostic angiography is excluded.

3) Benign Brain Tumour

2014

2019

Benign brain tumour means a nonmalignant tumour located in the cranial vault and limited to the brain, meninges or cranial nerves where all of the following conditions are met:

- It is life threatening:
- It has caused damage to the brain; It has undergone surgical removal or, if inoperable, has caused a permanent neurological deficit; and
- Its presence must be confirmed by a neurologist or neurosurgeon and supported by findings on Magnetic Resonance Imaging, Computerised Tomography, or other reliable imaging techniques.

The following are excluded:

- Cysts;
- Granulomas;
- Vascular Malformations;
- Haematomas; and
- Tumours of the pituitary gland or spinal cord.

Benign brain tumour means a nonmalignant tumour located in the cranial vault and limited to the brain, meninges or cranial nerves where all of the following conditions are met:

- It has undergone surgical removal or, if inoperable, has caused a permanent neurological deficit; and
- Its presence must be confirmed by a neurologist or neurosurgeon and supported by findings on Magnetic Resonance Imaging, Computerised Tomography, or other reliable imaging techniques.

The following are excluded:

- Cysts;
- Abscess:
- Angioma;
- · Granulomas;
- Vascular Malformations:
- Haematomas; and
- Tumours of the pituitary gland, spinal cord and skull base.

While there are less conditions to be met for 2019 definitions, there are more exclusion.

4) Blindness (Irreversible Loss of Sight)		
2014	2019	
Permanent and irreversible loss of sight in both eyes as a result of illness or Accident to the extent that even when tested with the use of visual aids, vision is measured at 3/60 or worse in both eyes using a Snellen eye chart or equivalent test, or visual field of twenty (20) degrees or less in both eyes.	Permanent and irreversible loss of sight in both eyes as a result of illness or accident to the extent that even when tested with the use of visual aids, vision is measured at 6/60 or worse in both eyes using a Snellen eye chart or equivalent test, or visual field of 20 degrees or less in both eyes. The blindness must be confirmed by an ophthalmologist.	
The blindness must be confirmed by an ophthalmologist	The blindness must not be correctable by surgical procedures, implants or any other means.	

The definition for 2019 is slightly better at 6/60 measurement of the vision.

·	
5) Coma	
2014	2019
A coma that persists for at least ninety-six (96) hours.	A coma that persists for at least 96 hours.
This diagnosis must be supported by evidence of all the following: (a) No response to external stimuli for at least ninety-six (96) hours; (b) Life support measures are necessary to sustain life; and (c) Brain damage resulting in permanent neurological deficit which must be assessed at least thirty (30) days after the onset of the coma.	This diagnosis must be supported by evidence of all of the following: • No response to external stimuli for at least 96 hours; • Life support measures are necessary to sustain life; and • Brain damage resulting in permanent neurological deficit which must be assessed at least 30 days after the onset of the coma.
Coma resulting directly from alcohol or drug abuse is excluded.	For the above definition, medically induced coma and coma resulting directly from alcohol or drug abuse are excluded.

6) Coronary Artery By-pass Surgery	
2014	2019
The actual undergoing of open-chest surgery or Minimally Invasive Direct Coronary Artery Bypass surgery to correct the narrowing or blockage of one or more coronary arteries with bypass grafts.	The actual undergoing of open-chest surgery or Minimally Invasive Direct Coronary Artery Bypass surgery to correct the narrowing or blockage of one or more coronary arteries with bypass grafts.
This diagnosis must be supported by angiographic evidence of significant coronary artery obstruction and the procedure must be considered medically necessary by a consultant cardiologist.	This diagnosis must be supported by angiographic evidence of significant coronary artery obstruction and the procedure must be considered medically necessary by a consultant cardiologist.
Angioplasty and all other intra-arterial, catheter-based techniques, 'keyhole' or laser procedures are excluded. No change	Angioplasty and all other intra-arterial, catheter-based techniques, 'keyhole' or laser procedures are excluded.

7) Deafness (Loss of Hearing)		
2014	2019	
Total and irreversible loss of hearing in both ears as a result of illness or Accident.	Total and irreversible loss of hearing in both ears as a result of illness or accident.	
This diagnosis must be supported by audiometric and sound-threshold tests provided and certified by an Ear, Nose, Throat (ENT) specialist. Total means "the loss of at least eighty (80) decibels in all frequencies of hearing".	This diagnosis must be supported by audiometric and sound-threshold tests provided and certified by an Ear, Nose, Throat (ENT) specialist. Total means "the loss of at least 80 decibels in all frequencies of hearing".	
	Irreversible means "cannot be reasonably restored to at least 40 decibels by medical treatment, hearing aid and/or surgical procedures consistent with the current standard of the medical services available in Singapore after a period of 6 months from the date of intervention."	
2019 definition is slightly more stringent with the mention of being irreversible		

8) End Stage Kidney Failure	
2014	2019
Chronic irreversible failure of both kidneys requiring either permanent renal dialysis or kidney transplantation.	Chronic irreversible failure of both kidneys requiring either permanent renal dialysis or kidney transplantation.
No change	

9) End Stage Liver Failure	
2014	2019
End stage liver failure as evidenced by all of the following: (a) Permanent jaundice; (b) Ascites; and (c) Hepatic encephalopathy.	End stage liver failure as evidenced by all of the following: • Permanent jaundice; • Ascites; and • Hepatic encephalopathy.
Liver disease secondary to alcohol or drug abuse is excluded.	Liver disease secondary to alcohol or drug abuse is excluded.
No change	

10) End Stage Lung Disease	
2014	2019
End stage lung disease, causing chronic respiratory failure.	End stage lung disease, causing chronic respiratory failure.
This diagnosis must be supported by evidence of all of the following: (a) FEV1 test results which are consistently less than one (1) litre; (b) Permanent supplementary oxygen therapy for hypoxemia; (c) Arterial blood gas analyses with partial oxygen pressures of 55mmHg or less (PaO2 ≤ 55mmHg); and (d) Dyspnea at rest.	This diagnosis must be supported by evidence of all of the following: • FEV1 test results which are consistently less than 1 litre; • Permanent supplementary oxygen therapy for hypoxemia; • Arterial blood gas analyses with partial oxygen pressures of 55mmHg or less (PaO2 ≤ 55mmHg); and • Dyspnea at rest.
The diagnosis must be confirmed by a respiratory physician.	The diagnosis must be confirmed by a respiratory physician.
No change	

11) Fulminant Hepatitis	
2014	2019
A submassive to massive necrosis of the liver by the Hepatitis virus, leading precipitously to liver failure.	A submassive to massive necrosis of the liver by the Hepatitis virus, leading precipitously to liver failure.
This diagnosis must be supported by all of the following: • Rapid decreasing of liver size as confirmed by abdominal ultrasound; • Necrosis involving entire lobules, leaving only a collapsed reticular framework; • Rapid deterioration of liver function tests; • Deepening jaundice; and • Hepatic encephalopathy.	This diagnosis must be supported by all of the following: Rapid decreasing of liver size as confirmed by abdominal ultrasound; Necrosis involving entire lobules, leaving only a collapsed reticular framework; Rapid deterioration of liver function tests; Deepening jaundice; and Hepatic encephalopathy.
No change	•

12) Heart Attack of Specified Severity

2014

2019

Death of heart muscle due to obstruction of blood flow, that is evident by at least three (3) of the following criteria proving the occurrence of a new heart attack:

- History of typical chest pain;
- New characteristic electrocardiographic changes; with the development of any of the following: ST elevation or depression, T wave inversion, pathological Q waves or left bundle
- Elevation of the cardiac biomarkers, inclusive of CKMB above the generally accepted normal laboratory levels or Cardiac Troponin T or I at 0.5ng/ml and above:
- Imaging evidence of new loss of viable myocardium or new regional wall motion abnormality.

The imaging must be done by a consultant cardiologist specified by the Company.

For the above definition, the following are excluded:

Angina;

branch block:

- Heart attack of indeterminate age; and
- A rise in cardiac biomarkers or Troponin T or I following an intraarterial cardiac procedure including, but not limited to, coronary angiography and coronary angioplasty.

Explanatory note: 0.5ng/ml = 0.5ug/L = 500pg/ml

Death of heart muscle due to ischaemia, that is evident by at least three of the following criteria proving the occurrence of a new heart attack:

- · History of typical chest pain;
- New characteristic electrocardiographic changes; with the development of any of the following: ST elevation or depression, T wave inversion, pathological Q waves or left bundle branch block:
- Elevation of the cardiac biomarkers, inclusive of CKMB above the generally accepted normal laboratory levels or Cardiac Troponin T or I at 0.5ng/ml and above:
- Imaging evidence of new loss of viable myocardium or new regional wall motion abnormality.

The imaging must be done by Cardiologist specified by the Company.

For the above definition, the following are excluded:

- · Angina;
- Heart attack of indeterminate age; and
- A rise in cardiac biomarkers or Troponin T or I following an intra-arterial cardiac procedure including, but not limited to, coronary angiography and coronary angioplasty.

Explanatory note: 0.5ng/ml = 0.5ug/L = 500pg/ml

No significant change

13) HIV Due to Blood Transfusion and Occupationally Acquired HIV

a) Infection with the Human Immunodeficiency Virus (HIV) through a blood transfusion, provided that all of the following conditions are met:

2014

- (i) The blood transfusion was medically necessary or given as part of a medical treatment:
- (ii) The blood transfusion was received in Singapore after the Policy Issue Date, issue date of this Supplementary Benefit, Benefit Commencement Date of this Supplementary Benefit, date of Endorsement or Reinstatement Date of this Supplementary Benefit, whichever is the later:
- (iii) The source of the infection is established to be from the Institution that provided the blood transfusion and the Institution is able to trace the origin of the HIV tainted blood; and
- (iv) The Life Assured does not suffer from Thalassaemia Major or Haemophilia.
- b) Infection with Human
 Immunodeficiency Virus
 (HIV) which resulted from an Accident
 occurring after the Policy Issue Date,
 issue date of this Supplementary Benefit,
 Benefit Commencement Date of this
 Supplementary Benefit, date of
 Endorsement or Reinstatement Date of
 this Supplementary Benefit, whichever is
 the later whilst the Life Assured was
 carrying out the normal professional
 duties of his or her occupation in
 Singapore, provided that all of the
 following are proven to Our satisfaction:
 (i) Proof of the Accident giving rise to the
- (i) Proof of the Accident giving rise to the infection must be reported to Us within thirty (30) days of the Accident taking place;
- (ii) Proof that the Accident involved a definite source of the HIV infected fluids;

A. Infection with the Human Immunodeficiency Virus (HIV) through a blood transfusion, provided that all of the following conditions are met:

2019

- The blood transfusion was medically necessary or given as part of a medical treatment:
- The blood transfusion was received in Singapore after the Issue Date, Date of endorsement or Date of reinstatement of this Supplementary Contract, whichever is the later; and
- The source of the infection is established to be from the Institution that provided the blood transfusion and the Institution is able to trace the origin of the HIV tainted blood.

- B. Infection with the Human Immunodeficiency Virus (HIV) which resulted from an accident occurring after the Issue Date, date of endorsement or date of reinstatement of this Supplementary Contract, whichever is the later whilst the Insured was carrying out the normal professional duties of his or her occupation in Singapore, provided that all of the following are proven to the Company's satisfaction:
- Proof that the accident involved a definite source of the HIV infected fluids;
- Proof of sero-conversion from HIV negative to HIV positive occurring during the 180 days after the documented

- (iii) Proof of sero-conversion from HIV negative to HIV positive occurring during the one hundred and eighty (180) days after the documented Accident. This proof must include a negative HIV antibody test conducted within five (5) days of the Accident; and
- (iv) HIV infection resulting from any other means including sexual activity and the use of intravenous drugs is excluded. This Clause 23B shall only be payable when the occupation of the Life Assured is a medical practitioner, houseman, medical student, state registered nurse, medical laboratory technician, dentist (surgeon and nurse) or paramedical worker, working in an approved medical centre or clinic (in Singapore).

This Supplementary Benefit shall not be payable by Us under a claim arising from Clause 23 in the event of a cure becoming available prior to the infection. "Cure" means any treatment that renders the HIV inactive or non-infectious.

- accident. This proof must include a negative HIV antibody test conducted within 5 days of the accident; and
- HIV infection resulting from any other means including sexual activity and the use of intravenous drugs is excluded.

This benefit is only payable when the occupation of the insured is a medical practitioner, housemen, medical student, state registered nurse, medical laboratory technician, dentist (surgeon and nurse) or paramedical worker, working in medical centre or clinic (in Singapore).

2019 definition is more favourable with less criteria to meet.

14) Idiopathic Parkinson's Disease

2014

2019

The unequivocal diagnosis of idiopathic Parkinson's Disease by a consultant neurologist. This diagnosis must be supported by all of the following conditions:

- The disease cannot be controlled with medication:
- There are signs of progressive impairment; and
- Inability of the Life Assured to perform (whether aided or unaided) at least three (3) of the six (6) "Activities of Daily Living" for a continuous period of at least six (6) months.

Activities of Daily Living:

- (i) Washing the ability to wash in the bath or shower (including getting into and out of the bath or shower) or wash satisfactorily by other means;
- (ii) Dressing the ability to put on, take off, secure and unfasten all garments and, as appropriate, any braces, artificial limbs or other surgical appliances;
- (iii) Transferring the ability to move from a bed to an upright chair or wheelchair and vice versa:
- (iv) Mobility the ability to move indoors from room to room on level surfaces:
- (v) Toileting the ability to use the lavatory or otherwise manage bowel and bladder functions so as to maintain a satisfactory level of personal hygiene;
- (vi) Feeding- the ability to feed oneself once food has been prepared and made available.

Drug-induced or toxic causes of Parkinsonism or all other causes of Parkinson's Disease are excluded.

For the purpose of this definition, "aided" shall mean with the aid of special equipment, device and/or apparatus and not pertaining to human aid.

The unequivocal diagnosis of idiopathic Parkinson's Disease by a consultant neurologist. This diagnosis must be supported by all of the following conditions:

- The disease cannot be controlled with medication; and
- Inability of the Life Assured to perform (whether aided or unaided) at least 3 of the 6 "Activities of Daily Living" for a continuous period of at least 6 months.

For the purpose of this definition, "aided" shall mean with the aid of special equipment, device and/or apparatus and not pertaining to human aid.

2019 definition is more favourable with less criteria to meet and no exclusion on drug-related or toxic causes.

15) Irreversible Aplastic Anaemia	
2014	2019
Chronic persistent bone marrow failure, confirmed by biopsy, which results in anaemia, neutropenia and thrombocytopenia requiring treatment with at least one (1) of the following:	Chronic persistent and irreversible bone marrow failure, confirmed by biopsy, which results in anaemia, neutropenia and thrombocytopenia requiring treatment with at least one of the following:
 Blood product transfusion; Marrow stimulating agents; Immunosuppressive agents; or Bone marrow transplantation. 	 Blood product transfusion; Bone marrow stimulating agents; Immunosuppressive agents; or Bone marrow or haematopoietic stem cell transplantation.
The diagnosis must be confirmed by a haematologist.	The diagnosis must be confirmed by a haematologist.
2014 definition is more favourable with less	criteria to meet but 2019 definition

16) **Irreversible Loss of Speech** 2014 2019 Total and irrecoverable loss of the ability Total and irreversible loss of the ability to speak as a result of injury or to speak as a result of injury or disease to the vocal cords. The inability to speak disease to the vocal cords. The must be established for a continuous inability to speak must be established period of twelve (12) months. This for a continuous period of 12 months. diagnosis must be supported by medical This diagnosis must be supported by evidence furnished by an Ear, Nose, medical evidence furnished by an Throat (ENT) specialist. Ear, Nose, Throat (ENT) specialist. All psychiatric related causes are All psychiatric related causes are excluded. excluded. No change

has a wide coverage of treatment

2014

2019

A condition as a result of a disease, illness or injury whereby the Life Assured is unable to perform (whether aided or unaided) at least three (3) of the six (6) "Activities of Daily Living", for a continuous period of six (6) months.

Activities of Daily Living:

- (i) Washing the ability to wash in the bath or shower (including getting into and out of the bath or shower) or wash satisfactorily by other means;
- (ii) Dressing the ability to put on, take off, secure and unfasten all garments and, as appropriate, any braces, artificial limbs or other surgical appliances;
- (iii) Transferring the ability to move from a bed to an upright chair or wheelchair and vice versa;
- (iv) Mobility the ability to move indoors from room to room on level surfaces;
- (v) Toileting the ability to use the lavatory or otherwise manage bowel and bladder functions so as to maintain a satisfactory level of personal hygiene;
- (vi) Feeding the ability to feed oneself once food has been prepared and made available.

This condition must be confirmed by the Company's approved doctor.
Non-organic diseases such as neurosis

and psychiatric illnesses are excluded. For the purpose of this definition, "aided" shall mean with the aid of special equipment, device and/or apparatus and not pertaining to human aid.

A condition as a result of a disease, illness or injury whereby the Life Assured is unable to perform (whether aided or unaided) at least 3 of the 6 "Activities of Daily Living", for a continuous period of 6 months.

This condition must be confirmed by the company's approved doctor. Non-organic diseases such as neurosis and psychiatric illnesses are excluded.

For the purpose of this definition, "aided" shall mean with the aid of special equipment, device and/or apparatus and not pertaining to human aid.

Others

The following two terms can be found in some of the above definitions, and their meanings are as follows:

1. Permanent Neurological DeficitPermanent means expected to last throughout the lifetime of the Life Assured.

Permanent neurological deficit means symptoms of dysfunction in the nervous system that are present on clinical examination and expected to last throughout the lifetime of the Life Assured. Symptoms that are covered include numbness, paralysis, localized weakness, dysarthria (difficulty with speech), aphasia (inability to speak), dysphagia (difficulty swallowing), visual impairment, difficulty in walking, lack of coordination, tremor, seizures, dementia, delirium and coma.

2. Activities of Daily Living (ADLs)

- (i) Washing the ability to wash in the bath or shower (including getting into and out of the bath or shower) or wash satisfactorily by other means;
- (ii) Dressing the ability to put on, take off, secure and unfasten all garments and, as appropriate, any braces,

	artificial limbs or other surgical
	appliances;
	(iii) Transferring - the ability to move
	from a bed to an upright chair or
	wheelchair and vice versa;
	(iv) Mobility - the ability to move
	indoors from room to room on level
	surfaces;
	(v) Toileting - the ability to use the
• 3	lavatory or otherwise manage bowel
	and bladder functions so as to maintain
	a satisfactory level of personal
	hygiene;
	(vi) Feeding - the ability to feed oneself
	once food has been prepared and
	made available.
No significant change	

18)	Major Burns	
	2014	2019
skin) burns percent (2	degree (full thickness of the scovering at least twenty 0%) of the surface Assured's body.	Third degree (full thickness of the skin) burns covering at least 20% of the surface of the Life Assured's body.
No change	Э	•

19) Major Cancers	
2014	2019
A malignant tumour positively diagnosed with histological confirmation and characterised by the uncontrolled growth of malignant cells with invasion and destruction of normal tissue.	A malignant tumour positively diagnosed with histological confirmation and characterised by the uncontrolled growth of malignant cells with invasion and destruction of normal tissue.
The term malignant tumour includes leukemia, lymphoma and sarcoma.	The term Major Cancer includes, but is not limited to, leukemia, lymphoma and sarcoma.
	Major Cancer diagnosed on the basis of finding tumour cells and/or tumour-associated molecules in blood, saliva, faeces, urine or any other bodily fluid in

For the above definition, the following are excluded:

- All tumours which are histologically classified as any of the following:
- Pre-malignant;
- Non-invasive;
- Carcinoma-in-situ;
- Having borderline malignancy;
- Having any degree of malignant potential;
- Having suspicious malignancy;
- Neoplasm of uncertain or unknown behavior;

or

- Cervical Dysplasia CIN-1, CIN-2 and CIN-3;
- Any non-melanoma skin carcinoma unless there is evidence of metastases to lymph nodes or beyond;
- Malignant melanoma that has not caused invasion beyond the epidermis;
- All Prostate cancers histologically described as T1N0M0 (TNM Classification) or below; or Prostate cancers of another equivalent or lesser classification:
- All Thyroid cancers histologically classified as T1N0M0 (TNM Classification) or below;
- All tumours of the Urinary Bladder histologically classified as T1N0M0 (TNM Classification) or below;
- All Gastro-Intestinal Stromal tumours histologically classified as T1N0M0 (TNM Classification) or below and with mitotic count of

less than or equal to 5/50 HPFs;

- Chronic Lymphocytic Leukaemia less than RAI Stage three (3); and
- All tumours in the presence of HIV infection.

the absence of further definitive and clinically verifiable evidence does not meet the above definition.

For the above definition, the following are excluded:

 All tumours which are histologically classified as any of the following:
 Pre-malignant;

Non-invasive:

Carcinoma-in-situ (Tis) or Ta; Having borderline malignancy; Having any degree of malignant potential:

Having suspicious malignancy; Neoplasm of uncertain or unknown behaviour: or

All grades of dysplasia, squamous intraepithelial lesions (HSIL and LSIL) and intra epithelial neoplasia;

- Any non-melanoma skin carcinoma, skin confined primary cutaneous lymphoma and dermatofibrosarcoma protuberans unless there is evidence of metastases to lymph nodes or beyond;
- Malignant melanoma that has not caused invasion beyond the epidermis;
- All Prostate cancers histologically described as T1N0M0 (TNM Classification) or below; or Prostate cancers of another equivalent or lesser classification;
- All Thyroid cancers histologically classified as T1N0M0 (TNM Classification) or below;
- All Neuroendocrine tumours histologically classified as T1N0M0 (TNM Classification) or below;
- All tumours of the Urinary Bladder histologically classified as T1N0M0 (TNM Classification) or below;
- All Gastro-Intestinal Stromal tumours histologically classified as Stage I or IA according to the latest edition of the AJCC Cancer Staging Manual, or below;

- Chronic Lymphocytic Leukaemia less than RAI Stage 3;
- All bone marrow malignancies which do not require recurrent blood transfusions, chemotherapy, targeted cancer therapies, bone marrow transplant, haematopoietic stem cell transplant or other major interventionist treatment; and
- All tumours in the presence of HIV infection.

2014 definition is more favourable with lesser exclusion.

2014

Accidental head injury resulting in permanentneurological deficit with persisting clinical symptoms to be assessed no sooner than six (6) weeks from the date of the Accident. This diagnosis must be confirmed by a consultant neurologist and supported by unequivocal findings on Magnetic Resonance Imaging, Computerised Tomography, or other reliable imaging techniques.

"Accident" means an event of violent, unexpected, external, involuntary and visible nature which is independent of any other cause and is the sole cause of the head Injury.

The following are excluded:

- Spinal cord injury; and
- Head injury due to any other causes.

Permanent means expected to last throughout the lifetime of the Life Assured.

Permanent neurological deficit with persisting clinical symptoms means symptoms of dysfunction in the nervous system that are present on clinical examination and expected to last throughout the lifetime of the Life Assured. Symptoms that are covered include numbness, paralysis, localised weakness, dysarthria (difficulty with speech), aphasia (inability to speak), dysphagia (difficulty swallowing), visual impairment, difficulty in walking, lack of coordination, tremor, seizures, dementia, delirium and coma.

Accidental head injury resulting in permanent neurological deficit to be assessed no sooner than 6 weeks from the date of the accident. This diagnosis must be confirmed by a consultant neurologist and supported by relevant findings on Magnetic Resonance Imaging, Computerised Tomography, or other reliable imaging techniques.

"Accident" means an event of violent, unexpected, external, involuntary and visible nature which is independent of any other cause and is the sole cause of the head Injury.

The following are excluded:

- · Spinal cord injury; and
- Head injury due to any other causes.

No significant changes

21) Major Organ / Bone Marrow Transplantation	
2014	2019
The receipt of a transplant of:	The receipt of a transplant of:
(a) Human bone marrow using	Human bone marrow using
haematopoietic stem cells preceded by	haematopoietic stem cells preceded
total bone marrow ablation; or (b) One (1)	by total bone marrow ablation; or
of the following human organs: heart,	One of the following human organs:
lung, liver, kidney, pancreas, that resulted	heart, lung, liver, kidney, pancreas,
from irreversible end stage failure of the	that resulted from irreversible end
relevant organ.	stage failure of the relevant organ.
Other stem cell transplants are excluded.	Other stem cell transplants are
	excluded.
No change	

22) Motor Neurone Disease	
2014	2019
Motor neurone disease characterised by progressive degeneration of corticospinal tracts and anterior horn cells or bulbar efferent neurones which include spinal muscular atrophy, progressive bulbar palsy, amyotrophic lateral sclerosis and primary lateral sclerosis.	Motor neurone disease characterised by progressive degeneration of corticospinal tracts and anterior horn cells or bulbar efferent neurones which include spinal muscular atrophy, progressive bulbar palsy, amyotrophic lateral sclerosis and primary lateral sclerosis.
This diagnosis must be confirmed by a neurologist as progressive and resulting in permanent neurological deficit.	This diagnosis must be confirmed by a neurologist as progressive and resulting in permanent neurological deficit.
No change	

23) Multiple Sclerosis	
2014	2019
The definite occurrence of Multiple Sclerosis. The diagnosis must be supported by all of the following: (a) Investigations which unequivocally confirm the diagnosis to be Multiple Sclerosis; (b) Multiple neurological deficits which occurred over a continuous period of at least six (6) months; and (c) Well documented history of exacerbations and remissions of said symptoms or neurological deficits.	The definite diagnosis of Multiple Sclerosis, and must be supported by all of the following: Investigations which unequivocally confirm the diagnosis to be Multiple Sclerosis; and Multiple neurological deficits which occurred over a continuous period of at least 6 months.
Other causes of neurological damage such as Systemic Lupus Erythematosus (SLE) with Lupus Nephritis and Human Immunodeficiency Virus (HIV) are excluded. 2019 definition is more favourable with less	Other causes of neurological damage such as SLE and HIV are excluded.

2019
The unequivocal diagnosis of muscular dystrophy must be made by a consultant neurologist.
6
The condition must result in the inability of the Life Assured to perform (whether aided or unaided) at least 3 of the 6 "Activities of Daily Living" for a continuous period of at least 6 months.
For the purpose of this definition, "aided" shall mean with the aid of special equipment, device and/or apparatus and not pertaining to human aid.

25) Open Chest Heart Valve Surgery	
2014	2019
The actual undergoing of open-heart surgery to replace or repair heart valve abnormalities.	The actual undergoing of open-heart surgery to replace or repair heart valve abnormalities.
The diagnosis of heart valve abnormality must be supported by cardiac catheterization or echocardiogram and the procedure must be considered medically necessary by a consultant cardiologist.	The diagnosis of heart valve abnormality must be supported by cardiac catheterization or echocardiogram and the procedure must be considered medically necessary by a consultant cardiologist.
No change	

26) Open Chest Surgery to Aorta	
2014	2019
The actual undergoing of major surgery to repair or correct an aneurysm, narrowing, obstruction or dissection of the Aorta through surgical opening of the chest or abdomen.	The actual undergoing of major surgery to repair or correct an aneurysm, narrowing, obstruction or dissection of the aorta through surgical opening of the chest or abdomen.
For the purpose of this definition, Aorta shall mean the thoracic and abdominal aorta but not its branches. Surgery performed using only minimally invasive or intra arterial techniques are excluded.	For the purpose of this definition, aorta shall mean the thoracic and abdominal aorta but not its branches. Surgery performed using only minimally invasive or intra-arterial techniques are excluded.
No change	

27) Other Serious Coronary Artery Disease	
2014	2019
The narrowing of the lumen of at least one (1) coronary artery by a minimum of seventy five percent (75%) and of two (2) others by a minimum of sixty percent (60%), as proven by coronary arteriography, regardless of whether or not any form of coronary artery surgery has been performed.	The narrowing of the lumen of at least one coronary artery by a minimum of 75% and of two others by a minimum of 60%, as proven by invasive coronary angiography, regardless of whether or not any form of coronary artery surgery has been performed.
	Diagnosis by Imaging or non-invasive diagnostic procedures such as CT scan or MRI does not meet the confirmatory status required by the definition.
Coronary arteries herein refer to left main stem, left anterior descending, circumflex and right coronary artery.	Coronary arteries herein refer to left main stem, left anterior descending, circumflex and right coronary artery. The branches of the above coronary arteries are excluded.
2014 definition is more favourable with less	ser criteria to meet

28) Paralysis (Loss of Use of Limbs)	
2014	2019
Total and irreversible loss of use of at least two (2) entire limbs due to injury or disease persisting for a period of at least six (6) weeks and with no foreseeable possibility of recovery.	Total and irreversible loss of use of at least 2 entire limbs due to injury or disease persisting for a period of at least 6 weeks and with no foreseeable possibility of recovery.
This condition must be confirmed by a consultant neurologist. Self-inflicted injuries are excluded.	This condition must be confirmed by a consultant neurologist. Self-inflicted injuries are excluded.
No change	

29) Persistent Vegetative State (Apallic Syndrome)	
2014	2019
Universal necrosis of the brain cortex with the brainstem intact. This diagnosis must be definitely confirmed by a consultant neurologist holding such an appointment at an approved hospital. This condition has to be medically documented for at least one (1) month.	Universal necrosis of the brain cortex with the brainstem intact. This diagnosis must be definitely confirmed by a consultant neurologist holding such an appointment at an approved hospital. This condition has to be medically documented for at least one month.
No change	

30) Poliomyelitis	
2014	2019
The occurrence of Poliomyelitis where the following conditions are met:	The occurrence of Poliomyelitis where the following conditions are met:
 Poliovirus is identified as the cause, Paralysis of the limb muscles or respiratory muscles must be present and persist for at least three (3) months. 	 Poliovirus is identified as the cause, Paralysis of the limb muscles or respiratory muscles must be present and persist for at least 3 months.
	The diagnosis must be confirmed by a consultant neurologist or specialist in the relevant medical field.
No significant change	

31) Primary Pulmonary Hyperte	31) Primary Pulmonary Hypertension	
2014	2019	
Primary Pulmonary Hypertension with substantial right ventricular enlargement confirmed by investigations including cardiac catheterisation, resulting in permanent physical impairment of at least Class IV of the New York Heart Association (NYHA) Classification of Cardiac Impairment.	Primary Pulmonary Hypertension with substantial right ventricular enlargement confirmed by investigations including cardiac catheterisation, resulting in permanent physical impairment of at least Class IV of the New York Heart Association (NYHA) Classification of Cardiac Impairment.	
The NYHA Classification of Cardiac Impairment:	The NYHA Classification of Cardiac Impairment:	
Class I: No limitation of physical activity. Ordinary physical activity does not cause undue fatigue, dyspnea, or anginal pain.	Class I: No limitation of physical activity. Ordinary physical activity does not cause undue fatigue, dyspnea, or anginal pain.	
Class II: Slight limitation of physical activity. Ordinary physical activity results in symptoms.	Class II: Slight limitation of physical activity. Ordinary physical activity results in symptoms.	
Class III: Marked limitation of physical activity. Comfortable at rest, but less than ordinary activity causes symptoms.	Class III: Marked limitation of physical activity. Comfortable at rest, but less than ordinary activity causes symptoms.	
Class IV: Unable to engage in any physical activity without discomfort. Symptoms may be present even at rest.	Class IV: Unable to engage in any physical activity without discomfort. Symptoms may be present even at rest.	
No change		

32) Progressive Scleroderma	
2014	2019
A systemic collagen-vascular disease causing progressive diffuse fibrosis in the skin, blood vessels and visceral organs.	A systemic collagen-vascular disease causing progressive diffuse fibrosis in the skin, blood vessels and visceral organs.
This diagnosis must be unequivocally supported by biopsy and serological evidence and the disorder must have reached systemic proportions to involve the heart, lungs or kidneys. The following are excluded:	This diagnosis must be unequivocally confirmed by a consultant rheumatologist and supported by biopsy or equivalent confirmatory test, and serological evidence, and the disorder must have reached systemic proportions to involve the heart, lungs or kidneys.
 Localised scleroderma (linear scleroderma or morphea); Eosinophilic fascitis; and CREST syndrome. 	The following are excluded: • Localised scleroderma (linear scleroderma or morphea); • Eosinophilic fascitis; and • CREST syndrome.
No significant change	

33) Severe Bacterial Meningitis		
2014	2019	
Bacterial infection resulting in severe inflammation of the membranes of the brain or spinal cord resulting in significant, irreversible and permanent neurological deficit. The neurological deficit must persist for at least six (6) weeks.	Bacterial infection resulting in severe inflammation of the membranes of the brain or spinal cord resulting in significant, irreversible and permanent neurological deficit. The neurological deficit must persist for at least 6 weeks.	
This diagnosis must be confirmed by: (a) The presence of bacterial infection in cerebrospinal fluid by lumbar puncture; and (b) A consultant neurologist. Bacterial Meningitis in the presence of HIV infection is excluded.	This diagnosis must be confirmed by: • The presence of bacterial infection in cerebrospinal fluid by lumbar puncture; and • A consultant neurologist. Bacterial Meningitis in the presence of HIV infection is excluded.	
No significant change		

34) Severe Encephalitis		
2014	2019	
Severe inflammation of brain substance (cerebral hemisphere, brainstem or cerebellum) caused by viral infection and resulting in permanent neurological deficit.	Severe inflammation of brain substance (cerebral hemisphere, brainstem or cerebellum) and resulting in permanent neurological deficit which must be documented for at least 6 weeks.	
This diagnosis must be certified by a consultant neurologist and the permanent neurological deficit must be documented for at least six (6) weeks.	This diagnosis must be certified by a consultant neurologist, and supported by any confirmatory diagnostic tests.	
Encephalitis caused by HIV infection is excluded.	Encephalitis caused by HIV infection is excluded.	
No significant change		

35) Stroke with Permanent Neurological Deficit

2014

2019

A cerebrovascular incident including infarction of brain tissue, cerebral and subarachnoid haemorrhage, intracerebral embolism and cerebral thrombosis resulting in permanent neurological deficit with persisting clinical symptoms. This diagnosis must be supported by all of the following conditions:

- Evidence of permanent clinical neurological deficit confirmed by a neurologist at least six (6) weeks after the event; and
- Findings on Magnetic Resonance Imaging, Computerised Tomography, or other reliable imaging techniques consistent with the diagnosis of a new stroke.

The following are excluded:

- Transient Ischaemic Attacks;
- Brain damage due to an Accident or injury, infection, vasculitis, and inflammatory disease;
- Vascular disease affecting the eye or optic nerve; and
- Ischaemic disorders of the vestibular system.

Permanent means expected to last throughout the lifetime of the Life Assured.

Permanent neurological deficit with persisting clinical symptoms means symptoms of dysfunction in the nervous system that are present on clinical examination and expected to last throughout the lifetime of the Life Assured.

Symptoms that are covered include numbness, paralysis, localised weakness, dysarthria (difficulty with speech), aphasia (inability to speak), dysphagia (difficulty swallowing), visual impairment, difficulty in walking, lack of coordination, tremor, seizures, dementia, delirium and coma. A cerebrovascular incident including infarction of brain tissue, cerebral and subarachnoid haemorrhage, intracerebral embolism and cerebral thrombosis resulting in permanent neurological deficit. This diagnosis must be supported by all of the following conditions:

- Evidence of permanent clinical neurological deficit confirmed by a neurologist at least 6 weeks after the event; and
- Findings on Magnetic Resonance Imaging, Computerised Tomography, or other reliable imaging techniques consistent with the diagnosis of a new stroke.

The following are excluded:

- Transient Ischaemic Attacks;
- Brain damage due to an accident or injury, infection, vasculitis, and inflammatory disease;
- Vascular disease affecting the eye or optic nerve;
- Ischaemic disorders of the vestibular system; and
- Secondary haemorrhage within a pre-existing cerebral lesion.

No significant change

36) Systemic Lupus Erythematosus with Lupus Nephritis

2014

2019

A multi-system, multifactorial, autoimmune disorder characterised by the development of auto-antibodies directed against various self-antigens. In respect of this contract, systemic lupus erythematosus will be restricted to those forms of systemic lupus erythematosus which involve the kidneys (Class III to Class V Lupus Nephritis, established by renal biopsy, and in accordance with the WHO Classification). The final diagnosis must be confirmed by a certified doctor specialising in Rheumatology and Immunology.

The unequivocal diagnosis of Systemic Lupus Erythematosus (SLE) based on recognised diagnostic criteria and supported with clinical and laboratory evidence. In respect of this contract, systemic lupus erythematosus will be restricted to those forms of systemic lupus erythematosus which involve the kidneys (Class III to Class VI Lupus Nephritis, established by renal biopsy, and in accordance with the RPS/ISN classification system). The final diagnosis must be confirmed by a certified doctor specialising in Rheumatology and Immunology.

The WHO Classification of Lupus Nephritis:

Class I Minimal Change Lupus Glomerulonephritis

Class II Messangial Lupus Glomerulonephritis

Class III Focal Segmental Proliferative Lupus Glomerulonephritis

Class IV Diffuse Proliferative Lupus Glomerulonephritis

Class V Membranous Lupus Glomerulonephritis The RPS/ISN Classification of lupus nephritis:

Class I Minimal mesangial lupus nephritis

Class II Mesangial proliferative lupus nephritis

Class III Focal lupus nephritis
(active and chronic;
proliferative and sclerosing)

Class IV Diffuse lupus nephritis (active and chronic; proliferative and sclerosing; segmental and global)

Class V Membranous lupus nephritis

Class VI Advanced sclerosis lupus nephritis

2019 definition has more classification of lupus nephritis

37) Terminal Illness	
2014	2019
The conclusive diagnosis of an illness	The conclusive diagnosis of an illness
that is expected to result in the death of	that is expected to result in the death of
the Life Assured within 12 months. This	the Life Assured within 12 months. This
diagnosis must be supported by a	diagnosis must be supported by a
specialist and confirmed by the	specialist and confirmed by the
Company's appointed doctor.	Company's appointed doctor.
61171.	
Terminal illness in the presence of HIV	Terminal illness in the presence of HIV
infection is excluded.	infection is excluded.
No significant change	